

Recombinant SARS-COV-2 Spike S1+S2 ECD(S-ECD) Protein

Catalog No.: RP01260LQ Recombinant 3 Publications

Sequence Information

Species Gene ID Swiss Prot SARS-COV-2 43740568

Tags

C-His

Synonyms

Envelope; SARS-CoV-2 Spike RBD (N501Y); Spike; Spike ECD; Spike RBD; Spike S1; Spike S2; Spike S2 ECD; S1-RBD protein; NCP-CoV RBD Protein; novel coronavirus RBD Protein; 2019-nCoV RBD Protein; Sglycoprotein Subunit1 RBD Protein

Product Information

Source PurificationHEK293 cells >97% by SDS-PAGE.

Endotoxin

< 0.1 EU/ μ g of the protein by LAL method.

Formulation

Supplied as a 0.22 µm filtered solution in PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Background

The spike protein (S) of coronavirus (CoV) attaches the virus to its cellular receptor, angiotensin-converting enzyme 2 (ACE2). A defined receptor-binding domain (RBD) on S mediates thisinteraction. The S protein plays key parts in the induction of neutralizing-antibody and T-cellresponses, as well as protective immunity.

Basic Information

Description

Recombinant SARS-COV-2 Spike S1+S2 ECD(S-ECD) Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Val11-Gln1208) of SARS-COV-2(2019-nCoV) S1+S2 ECD (S-ECD) (Accession #YP_009724390.1) fused with a 6×His tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Recombinant SARS-CoV-2 Spike S1+S2 ECD-His at 2 μ g/mL (100 μ L/well) can bind Recombinant Human ACE2 with a linear range of 0.15-9.92 ng/mL.

Storage


This product is stable at ≤ -70°C for up to 6 months from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature. Avoid repeated freeze-thaw cycles. Avoid repeated freeze/thaw cycles.

Contact


•

www.abclonal.com

Validation Data



Recombinant SARS-COV-2 S1+S2 ECD (S-ECD) Protein with His tag was determined by SDS-PAGE with Coomassie Blue, showing bands around 80,110,180 kDa.



Immobilized Recombinant SARS-COV-2 Spike S1+S2 ECD-His at $2\mu g/mL$ (100 $\mu L/well$) can bind Recombinant Human ACE2 with a linear range of 0.15-9.92 ng/mL.