### A0089

Leader in Biomolecular Solutions for Life Science

# p63 Mouse mAb

Catalog No.: A0089



# **Basic Information**

Observed MW 67kDa

Calculated MW 77kDa

Category SMab Recombinant Monoclonal Antibody

Applications WB,ELISA

Cross-Reactivity Human

# Background

This gene encodes a member of the p53 family of transcription factors. The functional domains of p53 family proteins include an N-terminal transactivation domain, a central DNAbinding domain and an oligomerization domain. Alternative splicing of this gene and the use of alternative promoters results in multiple transcript variants encoding different isoforms that vary in their functional properties. These isoforms function during skin development and maintenance, adult stem/progenitor cell regulation, heart development and premature aging. Some isoforms have been found to protect the germline by eliminating oocytes or testicular germ cells that have suffered DNA damage. Mutations in this gene are associated with ectodermal dysplasia, and cleft lip/palate syndrome 3 (EEC3); split-hand/foot malformation 4 (SHFM4); ankyloblepharon-ectodermal defects-cleft lip/palate; ADULT syndrome (acro-dermato-ungual-lacrimal-tooth); limb-mammary syndrome; Rap-Hodgkin syndrome (RHS); and orofacial cleft 8.

# **Recommended Dilutions**

### **Immunogen Information**

WB

# 1:500 - 1:1000

**Gene ID** 8626 Swiss Prot Q9H3D4

# Immunogen

A synthetic peptide of human p63

#### Synonyms

AIS; KET; LMS; NBP; RHS; p40; p51; p63; EEC3; OFC8; p73H; p73L; SHFM4; TP53L; TP73L; p53CP; TP53CP; B(p51A); B(p51B)

### Contact

# **Product Information**


 www.abclonal.com

Source Mouse **Isotype** IgG Purification Affinity purification

#### Storage

Store at 4°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using p63 Mouse mAb (A0089) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Mouse IgG (H+L) (AS003) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 5s.