

A0102

Leader in Biomolecular Solutions for Life Science



FOXO3A Rabbit pAb

Catalog No.: A0102

27 Publications

Basic Information

Observed MW

80-90kDa

Calculated MW

71kDa

Category

Polyclonal Antibody

Applications

WB, IHC-P, IF/ICC, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Enables DNA binding activity; DNA-binding transcription factor activity, RNA polymerase II-specific; and mitochondrial transcription factor activity. Involved in several processes, including mitochondrial transcription; positive regulation of muscle atrophy; and positive regulation of pri-miRNA transcription by RNA polymerase II. Acts upstream of or within several processes, including extrinsic apoptotic signaling pathway in absence of ligand; female gonad development; and neuronal stem cell population maintenance. Located in cytosol; mitochondrial outer membrane; and nucleus. Part of protein-containing complex. Colocalizes with mitochondrial matrix. Is expressed in several structures, including alimentary system; brain; early embryo; genitourinary system; and hemolymphoid system. Used to study dermoid cyst of ovary. Orthologous to human FOXO3 (forkhead box O3).

Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID

56484

Swiss Prot

Q9WVH4

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 558-635 of mouse FOXO3A (NP_062714.1).

Synonyms

Fkhr2; FKHL1; Foxo3a; 1110048B16Rik; 2010203A17Rik; FOXO3A

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.


Validation Data




Western blot analysis of extracts of various cell lines, using FOXO3A antibody (A0102) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 5s.




Immunohistochemistry analysis of paraffin-embedded Human colon using FOXO3A Rabbit pAb (A0102) at dilution of 1:50 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded Mouse kidney using FOXO3A Rabbit pAb (A0102) at dilution of 1:50 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of C6 cells using FOXO3A Rabbit pAb (A0102) at dilution of 100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH-3T3 cells using FOXO3A Rabbit pAb (A0102) at dilution of 100 (40x lens). Blue: DAPI for nuclear staining.