Leader in Biomolecular Solutions for Life Science

# MEK2 Rabbit pAb

Catalog No.: A0143



### **Basic Information**

### **Observed MW**

49kDa

#### **Calculated MW**

44kDa

#### Category

Polyclonal Antibody

### **Applications**

WB

#### **Cross-Reactivity**

Human

# **Background**

The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinase kinases. Mutations in this gene cause cardiofaciocutaneous syndrome (CFC syndrome), a disease characterized by heart defects, cognitive disability, and distinctive facial features similar to those found in Noonan syndrome. The inhibition or degradation of this kinase is also found to be involved in the pathogenesis of Yersinia and anthrax. A pseudogene, which is located on chromosome 7, has been identified for this gene.

# **Recommended Dilutions**

WB

1:500 - 1:1000

### **Immunogen Information**

Gene ID

Swiss Prot P36507

#### **Immunogen**

A synthetic peptide of human MEK2

### **Synonyms**

CFC4; MEK2; MKK2; MAPKK2; PRKMK2

### **Contact**

•

www.abclonal.com


## **Product Information**

**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at 4°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,pH7.3.

# **Validation Data**



Western blot analysis of lysates from HeLa cells, using MEK2 Rabbit pAb (A0143). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.