Leader in Biomolecular Solutions for Life Science

# SLC9A3R1 Rabbit pAb

Catalog No.: A0146 2 Publications



### **Basic Information**

### **Observed MW**

50kDa

### **Calculated MW**

39kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes a sodium/hydrogen exchanger regulatory cofactor. The protein interacts with and regulates various proteins including the cystic fibrosis transmembrane conductance regulator and G-protein coupled receptors such as the beta2-adrenergic receptor and the parathyroid hormone 1 receptor. The protein also interacts with proteins that function as linkers between integral membrane and cytoskeletal proteins. The protein localizes to actinrich structures including membrane ruffles, microvilli, and filopodia. Mutations in this gene result in hypophosphatemic nephrolithiasis/osteoporosis type 2, and loss of heterozygosity of this gene is implicated in breast cancer.

# **Recommended Dilutions**

WB

1:1000 - 1:4000

# **Immunogen Information**

Gene ID 9368 Swiss Prot

014745

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 189-358 of human SLC9A3R1 (NP 004243.1).

### **Synonyms**

EBP50; NHERF; NHE-RF; NHERF-1; NPHLOP2; SLC9A3R1

### **Contact**

•

www.abclonal.com

# **Product Information**

**Source** Rabbit **Isotype** IgG Purification


Affinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts of various cell lines, using SLC9A3R1 antibody (A0146) at 1:1000

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.