Leader in Biomolecular Solutions for Life Science

# ALDH1A1 Rabbit mAb

Catalog No.: A0157 Recombinant 2 Publications



# **Basic Information**

# **Observed MW**

55kDa

### **Calculated MW**

55kDa

### Category

SMab Recombinant Monoclonal Antibody

### **Applications**

WB,IF/ICC,IP,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

### CloneNo number

ARC52440

# **Background**

The protein encoded by this gene belongs to the aldehyde dehydrogenase family. Aldehyde dehydrogenase is the next enzyme after alcohol dehydrogenase in the major pathway of alcohol metabolism. There are two major aldehyde dehydrogenase isozymes in the liver, cytosolic and mitochondrial, which are encoded by distinct genes, and can be distinguished by their electrophoretic mobility, kinetic properties, and subcellular localization. This gene encodes the cytosolic isozyme. Studies in mice show that through its role in retinol metabolism, this gene may also be involved in the regulation of the metabolic responses to high-fat diet.

# **Recommended Dilutions**

**WB** 1:2000 - 1:20000

**IF/ICC** 1:50 - 1:200

IP 0.5μg-4μg antibody for 200μg-400μg extracts of

whole cells

# **Immunogen Information**

**Gene ID**216

Swiss Prot
P00352

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 250-350 of human ALDH1A1 (NP 000680.2).

# **Synonyms**

ALDC; ALDH1; HEL-9; HEL12; PUMB1; ALDH11; RALDH1; ALDH-E1; HEL-S-53e; ALDH1A1

### **Contact**

www.abclonal.com


# **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.



Western blot analysis of various lysates, using ALDH1A1 Rabbit mAb (A0157) at 1:20000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 10s.

Western blot analysis of various lysates, using ALDH1A1 Rabbit mAb (A0157) at 1:20000 dilution. Seconday analysis of HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.


Lysateš/počteieš: 25µg per lane.

100kDBlogking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit (RM00020).



Immunofluorescence analysis of A549 using ALDH1A1 Rabbit mAb (A0157) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunofluorescence analysis of Hep G2 using ALDH1A1 Rabbit mAb (A0157) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Immunoprecipitation analysis of 300  $\mu g$  extracts of A-549 cells using 3  $\mu g$  ALDH1A1 Rabbit mAb antibody (A0157). Western blot was performed from the immunoprecipitate using ALDH1A1 Rabbit mAb (A0157) at a dilition of 1:20000.