#### A0178

Leader in Biomolecular Solutions for Life Science

# FADS1 Rabbit mAb

Catalog No.: A0178 Recombinant



### **Basic Information**

**Observed MW** 52kDa

Calculated MW 52kDa

Category SMab Recombinant Monoclonal Antibody

Applications WB,ELISA

Cross-Reactivity Human, Mouse, Rat

CloneNo number ARC2499

### Background

The protein encoded by this gene is a member of the fatty acid desaturase (FADS) gene family. Desaturase enzymes regulate unsaturation of fatty acids through the introduction of double bonds between defined carbons of the fatty acyl chain. FADS family members are considered fusion products composed of an N-terminal cytochrome b5-like domain and a C-terminal multiple membrane-spanning desaturase portion, both of which are characterized by conserved histidine motifs. This gene is clustered with family members FADS1 and FADS2 at 11q12-q13.1; this cluster is thought to have arisen evolutionarily from gene duplication based on its similar exon/intron organization.

#### **Recommended Dilutions**

#### Immunogen Information

WB

1:500 - 1:1000

Gene ID 3992 **Swiss Prot** 060427

#### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 50-150 of human FADS1 (060427).

#### Synonyms

D5D; TU12; FADS6; FADSD5; LLCDL1; FADS1

Contact
---------

## **Product Information**

www.abclonal.com

**lsotype** IgG Purification Affinity purification


Storage

Source

Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of various lysates using FADS1 Rabbit mAb (A0178) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 1s.