Leader in Biomolecular Solutions for Life Science

# **DDIT3/CHOP Rabbit pAb**

Catalog No.: A0221 51 Publications



### **Basic Information**

### **Observed MW**

27kDa

### **Calculated MW**

19kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes a member of the CCAAT/enhancer-binding protein (C/EBP) family of transcription factors. The protein functions as a dominant-negative inhibitor by forming heterodimers with other C/EBP members, such as C/EBP and LAP (liver activator protein), and preventing their DNA binding activity. The protein is implicated in adipogenesis and erythropoiesis, is activated by endoplasmic reticulum stress, and promotes apoptosis. Fusion of this gene and FUS on chromosome 16 or EWSR1 on chromosome 22 induced by translocation generates chimeric proteins in myxoid liposarcomas or Ewing sarcoma. Multiple alternatively spliced transcript variants encoding two isoforms with different length have been identified.

# **Recommended Dilutions**

**WB** 1:1000 - 1:5000

IF/ICC 1:50 - 1:200

# **Immunogen Information**

Gene ID Swiss Prot 1649 P35638

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-169 of human DDIT3/CHOP (NP 004074.2).

### **Synonyms**

CHOP; CEBPZ; CHOP10; CHOP-10; GADD153; AltDDIT3; C/EBPzeta; DDIT3/CHOP

### **Contact**

www.abclonal.com


## **Product Information**

SourceIsotypePurificationRabbitIgGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of lysates from C6 cells, using DDIT3/CHOP Rabbit pAb (A0221) at 1:2000 dilution.C6 cells were treated by tunicamycin (2  $\mu$ g/ml) for 8 hours.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 30s.



Immunofluorescence analysis of L929 cells using DDIT3/CHOP Rabbit pAb (A0221) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of C6 cells using DDIT3/CHOP Rabbit pAb (A0221) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using DDIT3/CHOP Rabbit pAb (A0221) at dilution of 1:100. Blue: DAPI for nuclear staining.