

A0232

Leader in Biomolecular Solutions for Life Science



FABP4 Rabbit pAb

Catalog No.: A0232

3 Publications

Basic Information

Observed MW

14kDa

Calculated MW

15kDa

Category

Polyclonal Antibody

Applications

WB, IHC-P, IF/ICC, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

FABP4 encodes the fatty acid binding protein found in adipocytes. Fatty acid binding proteins are a family of small, highly conserved, cytoplasmic proteins that bind long-chain fatty acids and other hydrophobic ligands. It is thought that FABPs roles include fatty acid uptake, transport, and metabolism.

Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID

2167

Swiss Prot

P15090

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-132 of human FABP4 (NP_001433.1).

Synonyms

aP2; ALBP; AFABP; A-FABP; HEL-S-104; FABP4

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.



Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.

Validation Data




Western blot analysis of various lysates using FABP4 Rabbit pAb (A0232) at 1:1000 dilution.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 5s.

Western blot analysis of various lysates, using FABP4 Rabbit pAb (A0232) at 1:1000 dilution.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.
 Lysates/proteins: 25ug per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 5s.



Immunohistochemistry analysis of FABP4 in paraffin-embedded human breast tissue using FABP4 Rabbit pAb (A0232) at a dilution of 1:100 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunofluorescence analysis of paraffin-embedded Rat adipose using FABP4 Rabbit pAb (A0232) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (A5007) at 1:500 dilution. Blue: DAPI for nuclear staining.