Leader in Biomolecular Solutions for Life Science

# E-Cadherin Rabbit pAb

Catalog No.: A0314

**Basic Information** 

Mouse Monoclonal Antibody

**Observed MW** 

**Calculated MW** 

130kDa

97kDa

WB

Human

Category

Applications

**Cross-Reactivity** 



## Background

This gene encodes a classical cadherin of the cadherin superfamily. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate the mature glycoprotein. This calcium-dependent cell-cell adhesion protein is comprised of five extracellular cadherin repeats, a transmembrane region and a highly conserved cytoplasmic tail. Mutations in this gene are correlated with gastric, breast, colorectal, thyroid and ovarian cancer. Loss of function of this gene is thought to contribute to cancer progression by increasing proliferation, invasion, and/or metastasis. The ectodomain of this protein mediates bacterial adhesion to mammalian cells and the cytoplasmic domain is required for internalization. This gene is present in a gene cluster with other members of the cadherin family on chromosome 16.

## **Recommended Dilutions**

#### **Immunogen Information**

WB

# 1:500 - 1:2000

**Gene ID** 999 Swiss Prot P12830

**Immunogen** A synthetic peptide of human E-Cadherin

Synonyms UVO; CDHE; ECAD; LCAM; Arc-1; BCDS1; CD324; E-Cadherin

Contact

# Product Information


 www.abclonal.com

**Source** Rabbit **Isotype** IgG Purification Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# Validation Data



Western blot analysis of extracts of various cell lines, using E-Cadherin antibody (A0314). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.