#### A0380

Leader in Biomolecular Solutions for Life Science

1 Publications

# mTOR Rabbit pAb

Catalog No.: A0380

**Basic Information** 

Mouse Monoclonal Antibody

**Observed MW** 

**Calculated MW** 

280kDa

289kDa

Category

WB, IF/ICC

Human

Applications

**Cross-Reactivity** 



#### Background

The protein encoded by this gene belongs to a family of phosphatidylinositol kinase-related kinases. These kinases mediate cellular responses to stresses such as DNA damage and nutrient deprivation. This kinase is a component of two distinct complexes, mTORC1, which controls protein synthesis, cell growth and proliferation, and mTORC2, which is a regulator of the actin cytoskeleton, and promotes cell survival and cell cycle progression. This protein acts as the target for the cell-cycle arrest and immunosuppressive effects of the FKBP12-rapamycin complex. Inhibitors of mTOR are used in organ transplants as immunosuppressants, and are being evaluated for their therapeutic potential in SARS-CoV-2 infections. Mutations in this gene are associated with Smith-Kingsmore syndrome and somatic focal cortical dysplasia type II. The ANGPTL7 gene is located in an intron of this gene.

### **Recommended Dilutions**

WB	1:200 - 1:500
IF/ICC	1:50 - 1:200

### **Immunogen Information**

**Gene ID** 2475 Swiss Prot P42345

**Immunogen** A synthetic peptide of human mTOR

**Synonyms** SKS; FRAP; FRAP1; FRAP2; RAFT1; RAPT1; mTOR

### **Product Information**


 www.abclonal.com

**Source** Rabbit **Isotype** IgG Purification Affinity purification

Storage

Store at 4°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,pH7.3.

## Validation Data



Western blot analysis of extracts of various cell lines, using mTOR antibody (A0380). Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.