Leader in Biomolecular Solutions for Life Science

# N-Cadherin Rabbit pAb

Catalog No.: A0432 10 Publications



### **Basic Information**

### **Observed MW**

120kDa

### **Calculated MW**

100kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB,IHC-P,IF/ICC,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

### **Background**

This gene encodes a classical cadherin and member of the cadherin superfamily. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein is proteolytically processed to generate a calcium-dependent cell adhesion molecule and glycoprotein. This protein plays a role in the establishment of left-right asymmetry, development of the nervous system and the formation of cartilage and bone.

### **Recommended Dilutions**

WB	1:100 - 1:500
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

### **Immunogen Information**

Gene ID	<b>Swiss Prot</b>
1000	P19022

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 500-800 of human N-Cadherin (NP\_001783.2).

### **Synonyms**

CDHN; NCAD; ACOGS; ADHD8; CD325; ARVD14; CDw325; N-Cadherin

### **Contact**

<b>❸</b>	www.abclonal.com
----------	------------------

### **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of various lysates using N-Cadherin Rabbit pAb (A0432) at 1:500 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 1s.