

A0550

Leader in Biomolecular Solutions for Life Science



## Rig-I/DDX58 Rabbit pAb

Catalog No.: A0550

12 Publications

### Basic Information

#### Observed MW

107kDa

#### Calculated MW

107kDa

#### Category

Polyclonal Antibody

#### Applications

WB, ELISA

#### Cross-Reactivity

Human, Rat

### Background

DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases which are implicated in a number of cellular processes involving RNA binding and alteration of RNA secondary structure. This gene encodes a protein containing RNA helicase-DEAD box protein motifs and a caspase recruitment domain (CARD). It is involved in viral double-stranded (ds) RNA recognition and the regulation of the antiviral innate immune response. Mutations in this gene are associated with Singleton-Merten syndrome 2.

### Recommended Dilutions

WB 1:2000 - 1:4000

### Immunogen Information

#### Gene ID

23586

#### Swiss Prot

O95786

#### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 825-925 of human Rig-I/DDX58 (NP\_055129.2).

#### Synonyms

RIG1; DDX58; RIG-I; RLR-1; SGMRT2; Rig-I/DDX58

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification

Affinity purification


#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.

## Validation Data

---



Western blot analysis of lysates from HepG2 cells, using Rig-I/DDX58 Rabbit pAb (A0550) at 1:4000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.