

A0719

Leader in Biomolecular Solutions for Life Science



HMGB1 Rabbit pAb

Catalog No.: A0719

1 Publications

Basic Information

Observed MW

29kDa

Calculated MW

25kDa

Category

Mouse Monoclonal Antibody

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a protein that belongs to the High Mobility Group-box superfamily. The encoded non-histone, nuclear DNA-binding protein regulates transcription, and is involved in organization of DNA. This protein plays a role in several cellular processes, including inflammation, cell differentiation and tumor cell migration. Multiple pseudogenes of this gene have been identified. Alternative splicing results in multiple transcript variants that encode the same protein.

Recommended Dilutions

WB	1:100 - 1:500
IHC-P	1:50 - 1:100

Immunogen Information

Gene ID

3146

Swiss Prot

P09429

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 69-168 of human HMGB1 (NP_002119.1).

Synonyms

HMG1; HMG3; HMG-1; SBP-1; HMGB1

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH 7.3.


Validation Data




Western blot analysis of various lysates, using HMGB1 Rabbit pAb (A0719) at 1:400 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 20s.



Immunohistochemistry analysis of HMGB1 in paraffin-embedded human breast cancer using HMGB1 Rabbit pAb (A0719) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of HMGB1 in paraffin-embedded mouse testis using HMGB1 Rabbit pAb (A0719) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of HMGB1 in paraffin-embedded rat kidney using HMGB1 Rabbit pAb (A0719) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.