

A0743

Leader in Biomolecular Solutions for Life Science



## TEK Rabbit pAb

Catalog No.: A0743

2 Publications

### Basic Information

#### Observed MW

150kDa

#### Calculated MW

126kDa

#### Category

Polyclonal Antibody

#### Applications

WB, ELISA

#### Cross-Reactivity

Human, Mouse, Rat

### Background

This gene encodes a receptor that belongs to the protein tyrosine kinase Tie2 family. The encoded protein possesses a unique extracellular region that contains two immunoglobulin-like domains, three epidermal growth factor (EGF)-like domains and three fibronectin type III repeats. The ligand angiopoietin-1 binds to this receptor and mediates a signaling pathway that functions in embryonic vascular development. Mutations in this gene are associated with inherited venous malformations of the skin and mucous membranes. Alternative splicing results in multiple transcript variants. Additional alternatively spliced transcript variants of this gene have been described, but their full-length nature is not known.

### Recommended Dilutions

WB 1:100 - 1:500

### Immunogen Information

#### Gene ID

7010

#### Swiss Prot

Q02763

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 500-750 of human TEK (NP\_000450.2).

#### Synonyms

TIE2; VMCM; GLC3E; TIE-2; VMCM1; CD202B; TEK

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification

Affinity purification


#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.

## Validation Data

---



Western blot analysis of lysates from HUVEC cells, using TEK Rabbit pAb (A0743) at 1:500 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 90s.