Leader in Biomolecular Solutions for Life Science

www.abclonal.com

PKA C-alpha (PRKACA) Rabbit pAb

Catalog No.: A0798 1 Publications

Basic Information

Observed MW
40kDa

Calculated MW
41kDa

Category
Primary antibody

Applications
ELISA,WB

Cross-Reactivity
Human,Mouse,Rat

Background

This gene encodes one of the catalytic subunits of protein kinase A, which exists as a tetrameric holoenzyme with two regulatory subunits and two catalytic subunits, in its inactive form. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits have been identified in humans. cAMP-dependent phosphorylation of proteins by protein kinase A is important to many cellular processes, including differentiation, proliferation, and apoptosis. Constitutive activation of this gene caused either by somatic mutations, or genomic duplications of regions that include this gene, have been associated with hyperplasias and adenomas of the adrenal cortex and are linked to corticotropin-independent Cushing's syndrome. Alternative splicing results in multiple transcript variants encoding different isoforms. Tissue-specific isoforms that differ at the N-terminus have been described, and these isoforms may differ in the post-translational modifications that occur at the N-terminus of some isoforms.

Recommended Dilutions

WB	1:100 - 1:500

Immunogen Information

Gene ID
5566

Swiss Prot
P17612

Immunogen
A synthetic peptide corresponding to a sequence within amino acids 1-351 of human PKA C-alpha (PRKACA) (NP_002721.1).

Synonyms
CAFD1; PKACA; PPNAD4; PKA C-alpha (PRKACA)

Contact

www.abclonal.com

Product Information


Source
Rabbit

Isotype
IgG

Purification
Affinity purification

Storage
Store at -20℃. Avoid freeze / thaw cycles.
Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using PKA C-alpha (PRKACA) Rabbit pAb (A0798) at 1:500 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 10s.

For research use only. Not for therapeutic or diagnostic purposes.
Please visit http://abclonal.com for a complete listing of recommended products.