

A0818

Leader in Biomolecular Solutions for Life Science



## CD133 Rabbit mAb

Catalog No.: A0818

Recombinant

3 Publications

### Basic Information

#### Observed MW

120kDa

#### Calculated MW

97kDa

#### Category

SMab Recombinant Monoclonal Antibody

#### Applications

WB,IHC-P,ELISA

#### Cross-Reactivity

Human

#### CloneNo number

ARC2543

### Background

This gene encodes a pentaspan transmembrane glycoprotein. The protein localizes to membrane protrusions and is often expressed on adult stem cells, where it is thought to function in maintaining stem cell properties by suppressing differentiation. Mutations in this gene have been shown to result in retinitis pigmentosa and Stargardt disease. Expression of this gene is also associated with several types of cancer. This gene is expressed from at least five alternative promoters that are expressed in a tissue-dependent manner. Multiple transcript variants encoding different isoforms have been found for this gene.

### Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200

### Immunogen Information

#### Gene ID

8842

#### Swiss Prot

O43490

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 250-380 of human CD133 (O43490).

#### Synonyms

RP41; AC133; CD133; MCDR2; STGD4; CORD12; PROML1; MSTP061; 33

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification


Affinity purification

#### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,0.05% BSA,50% glycerol,pH7.3.


## Validation Data



Western blot analysis of lysates from HT-29 cells, using CD133 Rabbit mAb (A0818) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 1s.



Immunohistochemistry analysis of CD133 in paraffin-embedded human pancreas tissue using CD133 Rabbit mAb (A0818) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of CD133 in paraffin-embedded human kidney tissue using CD133 Rabbit mAb (A0818) at a dilution of 1:200 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.