Leader in Biomolecular Solutions for Life Science

# EIF4G1 Rabbit pAb

Catalog No.: A0881 2 Publications



### **Basic Information**

### **Observed MW**

250kDa

### **Calculated MW**

175kDa

#### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human

### **Background**

The protein encoded by this gene is a component of the multi-subunit protein complex EIF4F. This complex facilitates the recruitment of mRNA to the ribosome, which is a rate-limiting step during the initiation phase of protein synthesis. The recognition of the mRNA cap and the ATP-dependent unwinding of 5'-terminal secondary structure is catalyzed by factors in this complex. The subunit encoded by this gene is a large scaffolding protein that contains binding sites for other members of the EIF4F complex. A domain at its N-terminus can also interact with the poly(A)-binding protein, which may mediate the circularization of mRNA during translation. Alternative splicing results in multiple transcript variants, some of which are derived from alternative promoter usage.

### **Recommended Dilutions**

WB

1:500 - 1:1000

### **Immunogen Information**

Gene ID

Swiss Prot

1981

Q04637

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 550-650 of human EIF4G1 (NP 886553.3).

### **Synonyms**

P220; EIF4F; EIF4G; EIF4GI; PARK18; EIF-4G1; EIF4G1

### **Contact**

•

www.abclonal.com

### **Product Information**

**Source** Rabbit **Isotype** IgG Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of HeLa, using EIF4G1 Rabbit pAb (A0881) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 60s.