

# [KO Validated] DJ-1/PARK7 Rabbit pAb

Catalog No.: A0987 KO Validated 1 Publications

### **Basic Information**

## **Observed MW**

20kDa

# **Calculated MW**

20kDa

### Category

Primary antibody

### **Applications**

ELISA, WB, IF/ICC

### **Cross-Reactivity**

Human, Mouse

# **Background**

The product of this gene belongs to the peptidase C56 family of proteins. It acts as a positive regulator of androgen receptor-dependent transcription. It may also function as a redox-sensitive chaperone, as a sensor for oxidative stress, and it apparently protects neurons against oxidative stress and cell death. Defects in this gene are the cause of autosomal recessive early-onset Parkinson disease 7. Two transcript variants encoding the same protein have been identified for this gene.

# **Recommended Dilutions**

WB 1:500 - 1:2000

IF/ICC 1:50 - 1:200

# **Immunogen Information**

**Gene ID Swiss Prot** 11315 Q99497

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-189 of human DJ-1/PARK7 (NP\_009193.2).

### **Synonyms**

DJ1; DJ-1; GATD2; HEL-S-67p; K7

### **Contact**

• www.abclonal.com


### **Product Information**

**Purification** Source Isotype Rabbit IgG Affinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.



Western blot analysis of extracts from normal (control) and DJ-1/PARK7 knockout (KO) 293T cells,

using DJ-1/PARK7 antibody (A0987) at 1:1000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3s.




Western blot analysis of extracts of various cell lines, using DJ-1/PARK7 antibody (A0987) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.



Immunofluorescence analysis of HeLa cells using DJ-1/PARK7 antibody (A0987). Blue: DAPI for nuclear staining.