

A10214

Leader in Biomolecular Solutions for Life Science



Eph receptor B3 (EPHB3) Rabbit pAb

Catalog No.: A10214

Basic Information

Observed MW

110kDa

Calculated MW

110kDa

Category

Mouse Monoclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. The Eph family of receptors are divided into two groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family. This gene encodes a receptor for ephrin-B family members.

Recommended Dilutions

WB 1:500 - 1:2000

Immunogen Information

Gene ID

2049

Swiss Prot

P54753

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 779-998 of human Eph receptor B3 (EPHB3) (NP_004434.2).

Synonyms

EK2; ETK2; HEK2; TYRO6; Eph receptor B3 (EPHB3)

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of lysates from HeLa cells, using Eph receptor B3 (EPHB3) Rabbit pAb (A10214) at 1:3000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.