

A10251

Leader in Biomolecular Solutions for Life Science



NR2F2 Rabbit pAb

Catalog No.: A10251

2 Publications

Basic Information

Observed MW

47kDa

Calculated MW

46kDa

Category

Mouse Monoclonal Antibody

Applications

WB, IHC-P, IF/ICC, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a member of the steroid thyroid hormone superfamily of nuclear receptors. The encoded protein is a ligand inducible transcription factor that is involved in the regulation of many different genes. Alternate splicing results in multiple transcript variants.

Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID

7026

Swiss Prot

P24468

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human NR2F2 (NP_066285.1).

Synonyms

ARPI; ARP-1; CHTD4; NF-E3; SRXX5; SVP40; COUPTF2; COUPTFB; TFCOUP2; COUPTFII; NR2F2

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.


Validation Data




Western blot analysis of various lysates, using NR2F2 antibody (A10251) at 1:700 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 180s.




Western blot analysis of various lysates, using NR2F2 antibody (A10251) at 1:700 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90s.



Immunohistochemistry analysis of paraffin-embedded human kidney using NR2F2 Rabbit pAb (A10251) at dilution of 1:50 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.




Immunohistochemistry analysis of paraffin-embedded mouse liver using NR2F2 Rabbit pAb (A10251) at dilution of 1:50 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat liver using NR2F2 Rabbit pAb (A10251) at dilution of 1:50 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.

Validation Data



Immunofluorescence analysis of A431 cells using NR2F2 Rabbit pAb (A10251) at dilution of 1:50 (40x lens). Blue: DAPI for nuclear staining.