Leader in Biomolecular Solutions for Life Science

Rad23B Rabbit pAb

Catalog No.: A1034 4 Publications



Basic Information

Observed MW

58kDa

Calculated MW

43kDa

Category

Antibody

Applications

WB, ELISA

Cross-Reactivity

Human

Background

The protein encoded by this gene is one of two human homologs of Saccharomyces cerevisiae Rad23, a protein involved in the nucleotide excision repair (NER). This protein was found to be a component of the protein complex that specifically complements the NER defect of xeroderma pigmentosum group C (XP-c) cell extracts in vitro. This protein was also shown to interact with, and elevate the nucleotide excision activity of 3-methyladenine-DNA glycosylase (MPG), which suggested a role in DNA damage recognition in base excision repair. This protein contains an N-terminal ubiquitin-like domain, which was reported to interact with 26S proteasome, and thus this protein may be involved in the ubiquitin mediated proteolytic pathway in cells. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

Recommended Dilutions

WB

1:500 - 1:1000

Immunogen Information

Gene ID

Swiss Prot

P54727

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 100-200 of human Rad23B (NP_002865.1).

Synonyms

P58; HR23B; HHR23B; Rad23B

Contact

•

www.abclonal.com

Product Information

Source Rabbit **Isotype** IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from SK-OV-3 cells using Rad23B Rabbit pAb(A1034) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.