### A1035

Leader in Biomolecular Solutions for Life Science

# StAR Rabbit pAb

Catalog No.: A1035 12 Publications



### **Basic Information**

**Observed MW** 28kDa

**Calculated MW** 32kDa

Category Mouse Monoclonal Antibody

Applications WB, IP, ELISA

**Cross-Reactivity** Human, Mouse

### Background

The protein encoded by this gene plays a key role in the acute regulation of steroid hormone synthesis by enhancing the conversion of cholesterol into pregnenolone. This protein permits the cleavage of cholesterol into pregnenolone by mediating the transport of cholesterol from the outer mitochondrial membrane to the inner mitochondrial membrane. Mutations in this gene are a cause of congenital lipoid adrenal hyperplasia (CLAH), also called lipoid CAH. A pseudogene of this gene is located on chromosome 13.

### **Recommended Dilutions**

WB	1:100 - 1:500
IP	0.5µg-4µg antibody for 200µg-400µg extracts of whole cells

### **Immunogen Information**

Gene ID 6770

**Swiss Prot** P49675

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 64-285 of human StAR (NP\_000340.2).

## Synonyms

STARD1; StAR

### Contact

### **Product Information**

€

www.abclonal.com

Purification Affinity purification

#### Storage

Source


Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.

Isotype

lgG

### **Validation Data**



Western blot analysis of various lysates using StAR Rabbit pAb (A1035) at 1:400 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.

Immunoprecipitation agalysis of 300  $\mu$ g extracts of K-562 cells using 3  $\mu$ g StAR antibody (A1035). Western blot was performed from the immunoprecipitate using StAR antibody (A1035) at a dilution of 1:1000.

