Leader in Biomolecular Solutions for Life Science

# Cytokeratin 2e (KRT2) Rabbit pAb

Catalog No.: A10375



### **Basic Information**

## **Observed MW**

65kDa

### **Calculated MW**

65kDa

### Category

Mouse Monoclonal Antibody

#### **Applications**

WB, ELISA

### **Cross-Reactivity**

Human, Mouse

# **Background**

The protein encoded by this gene is a member of the keratin gene family. The type II cytokeratins consist of basic or neutral proteins which are arranged in pairs of heterotypic keratin chains coexpressed during differentiation of simple and stratified epithelial tissues. This type II cytokeratin is expressed largely in the upper spinous layer of epidermal keratinocytes and mutations in this gene have been associated with bullous congenital ichthyosiform erythroderma. The type II cytokeratins are clustered in a region of chromosome 12q12-q13.

# **Recommended Dilutions**

WB

1:500 - 1:2000

# Immunogen Information

**Gene ID** 

Swiss Prot P35908

3849

### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 1-220 of human Cytokeratin 2e (KRT2) (NP\_000414.2).

### **Synonyms**

K2e; KRTE; CK-2e; KRT2A; KRT2E; Cytokeratin 2e (KRT2)

### **Contact**

•

www.abclonal.com

## **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of extracts of various cell lines, using Cytokeratin 2e (KRT2) antibody (A10375) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.