# ABclonal www.abclonal.com

## **MEGF10 Rabbit pAb**

Catalog No.: A10508 1 Publications

## **Basic Information**

#### **Observed MW**

130kDa

## **Calculated MW**

122kDa

#### Category

Primary antibody

## **Applications**

ELISA,WB

## **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene encodes a member of the multiple epidermal growth factor-like domains protein family. The encoded protein plays a role in cell adhesion, motility and proliferation, and is a critical mediator of apoptotic cell phagocytosis as well as amyloid-beta peptide uptake in the brain. Expression of this gene may be associated with schizophrenia, and mutations in this gene are a cause of early-onset myopathy, areflexia, respiratory distress, and dysphagia (EMARDD) as well as congenital myopathy with minicores. Alternatively spliced transcript variants have been observed for this gene.

## **Recommended Dilutions**

WB

1:500 - 1:2000

## **Immunogen Information**

**Gene ID** 84466 **Swiss Prot** 

6 Q96KG7

## **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 26-160 of human MEGF10 (NP\_115822.1).

#### **Synonyms**

SR-F3; EMARDD; CMYP10A; CMYP10B; MEGF10

#### **Contact**

•

www.abclonal.com

## **Product Information**

**Source** Rabbit Isotype

IgG


**Purification** 

Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of U-251 MG, using MEGF10 Rabbit pAb (A10508) at 1:2000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.