Leader in Biomolecular Solutions for Life Science

# TNFR2/TNFRSF1B Rabbit pAb

Catalog No.: A1095 1 Publications



### **Basic Information**

#### **Observed MW**

75kDa

#### **Calculated MW**

48kDa

### Category

Polyclonal Antibody

#### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

### **Background**

The protein encoded by this gene is a member of the TNF-receptor superfamily. This protein and TNF-receptor 1 form a heterocomplex that mediates the recruitment of two antiapoptotic proteins, c-IAP1 and c-IAP2, which possess E3 ubiquitin ligase activity. The function of IAPs in TNF-receptor signalling is unknown, however, c-IAP1 is thought to potentiate TNF-induced apoptosis by the ubiquitination and degradation of TNF-receptor-associated factor 2, which mediates anti-apoptotic signals. Knockout studies in mice also suggest a role of this protein in protecting neurons from apoptosis by stimulating antioxidative pathways.

### **Recommended Dilutions**

WB

1:500 - 1:1000

### **Immunogen Information**

Gene ID

Swiss Prot P20333

**Immunogen** A synthetic peptide corresponding to a sequence within amino acids 295-377 of human TNFR2/TNFRSF1B (NP 001057.1).

### **Synonyms**

 $p^{7}5$ ; TBPII; TNFBR; TNFR2; CD120b; TNFR1B; TNFR80; TNF-R75;  $p^{7}5$ TNFR; TNF-R-II; TNFR2/TNFRSF1B

#### **Contact**

•

www.abclonal.com

### **Product Information**

**Source** Isotype Rabbit IgG


**Purification**Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of extracts of various cell lines, using TNFR2/TNFRSF1B antibody (A1095) at 1:1000

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.