

A11097

Leader in Biomolecular Solutions for Life Science



## Histone H2AX Rabbit pAb

Catalog No.: A11097

1 Publications

### Basic Information

#### Observed MW

15kDa

#### Calculated MW

15kDa

#### Category

Polyclonal Antibody

#### Applications

WB,IHC-P,IF/ICC,ELISA

#### Cross-Reactivity

Human,Mouse,Rat,Other (Wide Range Predicted)

### Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif.

### Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

### Immunogen Information

#### Gene ID

3014

#### Swiss Prot

P16104

#### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 50-143 of human Histone H2AX (NP\_002096.1).

#### Synonyms

H2A.X; H2A/X; H2AFX; Histone H2AX

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification


Affinity purification

#### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.


## Validation Data




Western blot analysis of various lysates using Histone H2AX Rabbit pAb (A11097) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.




Immunohistochemistry analysis of Histone H2AX in paraffin-embedded rat testis using Histone H2AX Rabbit pAb (A11097) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.




Immunohistochemistry analysis of Histone H2AX in paraffin-embedded human tonsil using Histone H2AX Rabbit pAb (A11097) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.




Immunohistochemistry analysis of Histone H2AX in paraffin-embedded mouse brain using Histone H2AX Rabbit pAb (A11097) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Confocal immunofluorescence analysis of HeLa cells using Histone H2AX Rabbit pAb (A11097) at dilution of 1:400. Blue: DAPI for nuclear staining.



Confocal immunofluorescence analysis of U-2 OS cells using Histone H2AX Rabbit pAb (A11097) at dilution of 1:400. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of U2OS cells using Histone H2AX Rabbit pAb (A11097) at dilution of 100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.