

A11274

Leader in Biomolecular Solutions for Life Science



[KO Validated] NADPH oxidase 4 (NOX4) Rabbit pAb

Catalog No.: A11274

KO Validated

22 Publications

Basic Information

Observed MW

65kDa

Calculated MW

67kDa

Category

Polyclonal Antibody

Applications

WB, IHC-P, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene encodes a member of the NOX-family of enzymes that functions as the catalytic subunit the NADPH oxidase complex. The encoded protein is localized to non-phagocytic cells where it acts as an oxygen sensor and catalyzes the reduction of molecular oxygen to various reactive oxygen species (ROS). The ROS generated by this protein have been implicated in numerous biological functions including signal transduction, cell differentiation and tumor cell growth. A pseudogene has been identified on the other arm of chromosome 11. Alternative splicing results in multiple transcript variants.

Recommended Dilutions

WB 1:500 - 1:1000

IHC-P 1:50 - 1:200

Immunogen Information

Gene ID

50507

Swiss Prot

Q9NPH5

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 328-578 of human NADPH oxidase 4 (NOX4) (NP_058627.1).

Synonyms

KOX; KOX-1; RENOX; 4)

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of lysates from 293T cells, using [KO Validated] NADPH oxidase 4 (NOX4) Rabbit pAb (A11274) at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 90s.




Western blot analysis of lysates from wild type (WT) and NADPH oxidase 4 (NOX4) knockout (KO) HeLa cells, using [KO Validated] NADPH oxidase 4 (NOX4) Rabbit pAb (A11274) at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
Lysates/proteins: 25µg per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit (RM00020).
Exposure time: 60s.



Immunohistochemistry analysis of NADPH oxidase 4 (NOX4) in paraffin-embedded human kidney using [KO Validated] NADPH oxidase 4 (NOX4) Rabbit pAb (A11274) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of NADPH oxidase 4 (NOX4) in paraffin-embedded mouse colon using [KO Validated] NADPH oxidase 4 (NOX4) Rabbit pAb (A11274) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of NADPH oxidase 4 (NOX4) in paraffin-embedded rat colon using [KO Validated] NADPH oxidase 4 (NOX4) Rabbit pAb (A11274) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.