A11777

Leader in Biomolecular Solutions for Life Science

Catalase Rabbit pAb

Catalog No.: A11777 13 Publications



Basic Information

Observed MW 60kDa

Calculated MW 60kDa

Category Polyclonal Antibody

Applications WB,IHC-P,IF/ICC,ELISA

Cross-Reactivity Human,Mouse,Rat

Background

This gene encodes catalase, a key antioxidant enzyme in the bodies defense against oxidative stress. Catalase is a heme enzyme that is present in the peroxisome of nearly all aerobic cells. Catalase converts the reactive oxygen species hydrogen peroxide to water and oxygen and thereby mitigates the toxic effects of hydrogen peroxide. Oxidative stress is hypothesized to play a role in the development of many chronic or late-onset diseases such as diabetes, asthma, Alzheimer's disease, systemic lupus erythematosus, rheumatoid arthritis, and cancers. Polymorphisms in this gene have been associated with decreases in catalase activity but, to date, acatalasemia is the only disease known to be caused by this gene.

Recommended Dilutions

Immunogen Information

WB	1:500 - 1:2000	Gene ID 847	Swiss Prot P04040
IHC-P	1:50 - 1:100		
IF/ICC	1:50 - 1:100	Immunogen Recombinant fusion protein containing a sequence corresponding to amino acids 1-225 of human Catalasealase (NP 001743.1).	

Synonyms

CAT; catalase; MGC138422; MGC138424; Catalase

Product Information

www.abclonal.com

lsotype lgG Purification Affinity purification


Storage

Source


Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of various lysates using Catalase Rabbit pAb (A11777) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 5s.



Immunofluorescence analysis of HeLa cells using Catalase Rabbit pAb (A11777) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.