

A12296

Leader in Biomolecular Solutions for Life Science



## RNF10 Rabbit pAb

Catalog No.: A12296

### Basic Information

**Observed MW**

90kDa

**Calculated MW**

90kDa

**Category**

Polyclonal Antibody

**Applications**

WB, ELISA

**Cross-Reactivity**

Mouse, Rat

### Background

The protein encoded by this gene contains a ring finger motif, which is known to be involved in protein-protein interactions. The specific function of this protein has not yet been determined. EST data suggests the existence of multiple alternatively spliced transcript variants, however, their full length nature is not known.

### Recommended Dilutions

WB 1:100 - 1:500

### Immunogen Information

**Gene ID**

9921

**Swiss Prot**

Q8N5U6

**Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 552-811 of human RNF10 (NP\_055683.3).

**Synonyms**

RIE2; RNF10

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

**Source**

Rabbit

**Isotype**

IgG

**Purification**


Affinity purification

**Storage**

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3.

## Validation Data



Western blot analysis of lysates from Rat testis, using RNF10 Rabbit pAb (A12296) at 1:400 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 90s.

Western blot analysis of lysates from Mouse testis using RNF10 Rabbit pAb(A12296) at 1:500 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L)(AS014) at 1:10000 dilution.  
Lysates/proteins: 25 µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 10s.

