### A12309

Leader in Biomolecular Solutions for Life Science

# NOX1 Rabbit pAb

Catalog No.: A12309 4 Publications



### **Basic Information**

Observed MW 65kDa

Calculated MW 65kDa

Category Polyclonal Antibody

Applications WB,IF/ICC,ELISA

Cross-Reactivity Human

### Background

This gene encodes a member of the NADPH oxidase family of enzymes responsible for the catalytic one-electron transfer of oxygen to generate superoxide or hydrogen peroxide. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene.

## **Recommended Dilutions**

### Immunogen Information

1:100 - 1:500	Gene ID	Swiss Prot
	27035	Q9Y5S8
1:50 - 1:200		

#### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 200-300 of human NOX1 (NP\_008983.2).

### Synonyms

MOX1; NOH1; NOH-1; GP91-2; NOH-1L; NOX1

WB

**IF/ICC** 

### **Product Information**

www.abclonal.com

Purification Affinity purification

### Storage

Source


Rabbit

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.


Isotype

lgG


# **Validation Data**



Western blot analysis of various lysates using NOX1 Rabbit pAb (A12309) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 10s.



Immunofluorescence analysis of HepG2 cells using NOX1 Rabbit pAb (A12309) at dilution of 1:50 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HT-29 cells using NOX1 Rabbit pAb (A12309) at dilution of 1:50 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.