

PCNA Rabbit mAb

Catalog No.: A12427 **Recombinant** **34 Publications**

Basic Information

Observed MW

35kDa

Calculated MW

29kDa

Category

Primary antibody

Applications

ELISA, WB, IHC-P

Cross-Reactivity

Human, Mouse, Rat, African green monkey

CloneNo number

ARC51324

Recommended Dilutions

WB 1:500 - 1:1000

IHC-P 1:2000 - 1:10000

Background

The protein encoded by this gene is found in the nucleus and is a cofactor of DNA polymerase delta. The encoded protein acts as a homotrimer and helps increase the processivity of leading strand synthesis during DNA replication. In response to DNA damage, this protein is ubiquitinated and is involved in the RAD6-dependent DNA repair pathway. Two transcript variants encoding the same protein have been found for this gene. Pseudogenes of this gene have been described on chromosome 4 and on the X chromosome.

Immunogen Information

Gene ID

5111

Swiss Prot

P12004

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 162-261 of human PCNA (NP_002583.1).

Synonyms

ATLD2; PCNA

Contact

 | www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 0.05% BSA, 50% glycerol, pH7.3.


Validation Data




Western blot analysis of various lysates, using PCNA Rabbit mAb (A12427) at 1:1000 dilution.
 Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.
 Lysates/proteins: 25µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit (RM00020).
 Exposure time: 30s.




Immunohistochemistry analysis of PCNA in paraffin-embedded human colon carcinoma tissue using PCNA Rabbit mAb (A12427) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of PCNA in paraffin-embedded mouse testis tissue using PCNA Rabbit mAb (A12427) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of PCNA in paraffin-embedded human esophageal cancer tissue using PCNA Rabbit mAb (A12427) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of PCNA in paraffin-embedded rat testis tissue using PCNA Rabbit mAb (A12427) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of PCNA in paraffin-embedded human esophagus tissue using PCNA Rabbit mAb (A12427) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.




Immunohistochemistry analysis of PCNA in paraffin-embedded human colon tissue using PCNA Rabbit mAb (A12427) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.

Validation Data



Immunohistochemistry analysis of PCNA in paraffin-embedded rat colon tissue using PCNA Rabbit mAb (A12427) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of PCNA in paraffin-embedded mouse testis tissue using PCNA Rabbit mAb (A12427) at a dilution of 1:8000 (40x lens). High pressure antigen retrieval was performed with 0.01 M citrate buffer (pH 6.0) prior to IHC staining.