

A12814

Leader in Biomolecular Solutions for Life Science



## Olig2 Rabbit pAb

Catalog No.: A12814

### Basic Information

#### Observed MW

35kDa

#### Calculated MW

32kDa

#### Category

Mouse Monoclonal Antibody

#### Applications

WB, IHC-P, ELISA

#### Cross-Reactivity

Human, Mouse, Rat

### Background

This gene encodes a basic helix-loop-helix transcription factor which is expressed in oligodendroglial tumors of the brain. The protein is an essential regulator of ventral neuroectodermal progenitor cell fate. The gene is involved in a chromosomal translocation t(14;21)(q11.2;q22) associated with T-cell acute lymphoblastic leukemia. Its chromosomal location is within a region of chromosome 21 which has been suggested to play a role in learning deficits associated with Down syndrome.

### Recommended Dilutions

WB	1:500 - 1:1000
IHC-P	1:50 - 1:100

### Immunogen Information

#### Gene ID

10215

#### Swiss Prot

Q13516

#### Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-80 of human Olig2 (NP\_005797.1).

#### Synonyms

BHLHB1; OLIGO2; RACK17; PRKCBP2; bHLHe19; Olig2

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification


Affinity purification

#### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal, 50% glycerol, pH7.3.


## Validation Data



Western blot analysis of various lysates using Olig2 Rabbit pAb (A12814) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 5s.



Immunohistochemistry analysis of Olig2 in paraffin-embedded rat brain using Olig2 Rabbit pAb (A12814) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of Olig2 in paraffin-embedded mouse brain using Olig2 Rabbit pAb (A12814) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.