Leader in Biomolecular Solutions for Life Science

# **DRD2** Rabbit pAb

Catalog No.: A12930 5 Publications



#### **Basic Information**

#### **Observed MW**

51kDa

#### **Calculated MW**

51kDa

#### Category

Polyclonal Antibody

### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

This gene encodes the D2 subtype of the dopamine receptor. This G-protein coupled receptor inhibits adenylyl cyclase activity. A missense mutation in this gene causes myoclonus dystonia; other mutations have been associated with schizophrenia. Alternative splicing of this gene results in two transcript variants encoding different isoforms. A third variant has been described, but it has not been determined whether this form is normal or due to aberrant splicing.

# **Recommended Dilutions**

WB

1:100 - 1:500

# **Immunogen Information**

Gene ID 1813 Swiss Prot P14416

# Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human DRD2 (NP 000786.1).

### **Synonyms**

D2R; D2DR; DRD2

### **Contact**

•

www.abclonal.com


## **Product Information**

**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.



Western blot analysis of U-251MG, using DRD2 antibody (A12930) at 1:400 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.