Leader in Biomolecular Solutions for Life Science

# **CEL Rabbit pAb**

Catalog No.: A13011



# **Basic Information**

### **Observed MW**

67kDa

#### **Calculated MW**

79kDa

### Category

Mouse Monoclonal Antibody

#### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Mouse

# **Background**

The protein encoded by this gene is a glycoprotein secreted from the pancreas into the digestive tract and from the lactating mammary gland into human milk. The physiological role of this protein is in cholesterol and lipid-soluble vitamin ester hydrolysis and absorption. This encoded protein promotes large chylomicron production in the intestine. Also its presence in plasma suggests its interactions with cholesterol and oxidized lipoproteins to modulate the progression of atherosclerosis. In pancreatic tumoral cells, this encoded protein is thought to be sequestrated within the Golgi compartment and is probably not secreted. This gene contains a variable number of tandem repeat (VNTR) polymorphism in the coding region that may influence the function of the encoded protein.

# **Recommended Dilutions**

WB

1:500 - 1:2000

## **Immunogen Information**

Gene ID

**Swiss Prot** 

P19835

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 350-550 of human CEL (NP 001798.2).

### **Synonyms**

BAL; FAP; BSDL; BSSL; CELL; FAPP; LIPA; CEase; MODY8; CEL

### **Contact**

•

www.abclonal.com

## **Product Information**

**Source** Rabbit **Isotype** IgG

Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of lysates from mouse pancreas, using CEL Rabbit pAb (A13011) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.