A13336

Leader in Biomolecular Solutions for Life Science

PCNA Rabbit pAb

Catalog No.: A13336 7 Publications



Basic Information

Observed MW 34kDa

Calculated MW 29kDa

Category Polyclonal Antibody

Applications WB,IHC-P,ELISA

Cross-Reactivity Human,Mouse,Rat

Background

The protein encoded by this gene is found in the nucleus and is a cofactor of DNA polymerase delta. The encoded protein acts as a homotrimer and helps increase the processivity of leading strand synthesis during DNA replication. In response to DNA damage, this protein is ubiquitinated and is involved in the RAD6-dependent DNA repair pathway. Two transcript variants encoding the same protein have been found for this gene. Pseudogenes of this gene have been described on chromosome 4 and on the X chromosome.

Recommended Dilutions

Immunogen Information

1:500 - 1:1000	Gene ID	Swiss Prot
	5111	P12004
1:50 - 1:200		

Immunogen

A synthetic peptide corresponding to a sequence within a mino acids 162-261 of human PCNA (NP_002583.1).

Synonyms ATLD2; PCNA

WB

IHC-P

Product Information


 www.abclonal.com

Source Rabbit **Isotype** IgG Purification Affinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data




Western blot analysis of extracts of various cell lines, using PCNA antibody (A13336) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. BioCRing buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 3s. 45kDa 25kDa 25kDa




15kDa

Immunohistochemistry analysis of paraffinembedded mouse colon using PCNA Rabbit pAb (A13336) at dilution of 1:200 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffinembedded mouse testis using PCNA Rabbit pAb (A13336) at dilution of 1:200 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffinembedded rat testis using PCNA Rabbit pAb (A13336) at dilution of 1:200 (40x lens).Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.