A13670

Leader in Biomolecular Solutions for Life Science

UBE2L6 Rabbit pAb

Catalog No.: A13670



Basic Information

Observed MW 18kDa

Calculated MW 18kDa

Category Polyclonal Antibody

Applications WB,IF/ICC,ELISA

Cross-Reactivity Human,Mouse,Rat

Background

The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes (E1s), ubiquitin-conjugating enzymes (E2s) and ubiquitin-protein ligases (E3s). This gene encodes a member of the E2 ubiquitin-conjugating enzyme family. This enzyme is highly similar in primary structure to the enzyme encoded by the UBE2L3 gene. Two alternatively spliced transcript variants encoding distinct isoforms have been found for this gene.

Recommended Dilutions

Immunogen Information

WB	1:100 - 1:500	Gene
		9246
IF/ICC	1:50 - 1:200	

www.abclonal.com

0	Gene ID	Swiss Prot	
	9246	014933	
n			

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 100-153 of human UBE2L6 (NP_004214.1).

Synonyms RIG-B; UBCH8; UBE2L6

€


Product Information

Source Rabbit **lsotype** IgG Purification Affinity purification


Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.


Validation Data



Western blot analysis of various lysates, using UBE2L6 antibody (A13670) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.



Immunofluorescence analysis of HeLa cells using UBE2L6 antibody (A13670) at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of NIH/3T3 cells using UBE2L6 antibody (A13670) at dilution of 1:100. Blue: DAPI for nuclear staining.