

# Coagulation Protein C Rabbit pAb

Catalog No.: A1432

## Basic Information

### Observed MW

50kDa

### Calculated MW

52kDa

### Category

Primary antibody

### Applications

ELISA, WB

### Cross-Reactivity

Human, Mouse

## Background

This gene encodes a vitamin K-dependent plasma glycoprotein. The encoded protein is cleaved to its activated form by the thrombin-thrombomodulin complex. This activated form contains a serine protease domain and functions in degradation of the activated forms of coagulation factors V and VIII. Mutations in this gene have been associated with thrombophilia due to protein C deficiency, neonatal purpura fulminans, and recurrent venous thrombosis.

## Recommended Dilutions

WB 1:500 - 1:2000

## Immunogen Information

### Gene ID

5624

### Swiss Prot

P04070

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 43-300 of human Coagulation Protein C (NP\_000303.1).

### Synonyms

PC; APC; PROC1; THPH3; THPH4; Coagulation Protein C

## Contact

 | [www.abclonal.com](http://www.abclonal.com)

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Validation Data



Western blot analysis of extracts of ES-2 cells, using Coagulation Protein C antibody (A1432) at 1:1000 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (A5014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.