Leader in Biomolecular Solutions for Life Science

# **CCR9 Rabbit pAb**

Catalog No.: A14848 3 Publications



### **Basic Information**

### **Observed MW**

42kDa

#### **Calculated MW**

42kDa

#### Category

Mouse Monoclonal Antibody

#### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

### **Background**

The protein encoded by this gene is a G protein-coupled receptor with seven transmembrane domains that belongs to the beta chemokine receptor family. Chemokines and their receptors are key regulators of thymocyte migration and maturation in normal and inflammation conditions. This gene is differentially expressed in T lymphocytes of the small intestine and colon, and its interaction with chemokine 25 contributes to intestinal intraepithelial lymphocyte homing to the small intestine. This suggests a role for this gene in directing immune responses to different segments of the gastrointestinal tract. This gene and its exclusive ligand, chemokine 25, are overexpressed in a variety of malignant tumors and are closely associated with tumor proliferation, apoptosis, invasion, migration and drug resistance. This gene maps to the chemokine receptor gene cluster. Multiple transcript variants encoding different isoforms have been found for this gene.

### **Recommended Dilutions**

WB

1:500 - 1:2000

### **Immunogen Information**

**Gene ID** 10803

Swiss Prot

P51686

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 300-369 of human CCR9 (NP 112477.1).

### **Synonyms**

GPR28; CDw199; GPR-9-6; CC-CKR-9; CCR9

### **Contact**

•

www.abclonal.com

### **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## **Validation Data**



Western blot analysis of lysates from Mouse brain, using CCR9 Rabbit pAb (A14848) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.