

BAIAP2L1 Rabbit pAb

Catalog No.: A15173

Basic Information

Observed MW

Calculated MW

57kDa

Category

Primary antibody

Applications

ELISA, IF/ICC

Cross-Reactivity

Human

Background

This gene encodes a member of the IMD (IRSp53/MIM homology domain) family. Members of this family can be subdivided in two groups, the IRSp53-like and MIM-like, based on the presence or absence of the SH3 (Src homology 3) domain. The protein encoded by this gene contains a conserved IMD, also known as F-actin bundling domain, at the N-terminus, and a canonical SH3 domain near the C-terminus, so it belongs to the IRSp53-like group. This protein is the substrate for insulin receptor tyrosine kinase and binds to the small GTPase Rac. It is involved in signal transduction pathways that link deformation of the plasma membrane and remodeling of the actin cytoskeleton. It also promotes actin assembly and membrane protrusions when overexpressed in mammalian cells, and is essential to the formation of a potent actin assembly complex during EHEC (Enterohemorrhagic Escherichia coli) pedestal formation.

Recommended Dilutions

IF/ICC 1:50 - 1:200

Immunogen Information

Gene ID

55971

Swiss Prot

Q9UHR4

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human BAIAP2L1 (NP_061330.2).

Synonyms

IRTKS; BAIAP2L1

Contact

 | www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Validation Data



Immunofluorescence analysis of U2OS cells using BAIAP2L1 Rabbit pAb (A15173) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.