### A15181

Leader in Biomolecular Solutions for Life Science

# MCCC2 Rabbit pAb

Catalog No.: A15181



## **Basic Information**

**Observed MW** 61kDa

**Calculated MW** 61kDa

Category Polyclonal Antibody

Applications WB,IHC-P,IF/ICC,ELISA

**Cross-Reactivity** Human, Mouse, Rat

## Background

This gene encodes the small subunit of 3-methylcrotonyl-CoA carboxylase. This enzyme functions as a heterodimer and catalyzes the carboxylation of 3-methylcrotonyl-CoA to form 3-methylglutaconyl-CoA. Mutations in this gene are associated with 3-Methylcrotonylglycinuria, an autosomal recessive disorder of leucine catabolism. Alternative splicing results in multiple transcript variants encoding distinct isoforms.

## **Recommended Dilutions**

### **Immunogen Information**

| 1:500 - 1:2000 | Gene ID | Swiss Prot |  |
|----------------|---------|------------|--|
|                | 64087   | Q9HCC0     |  |
| 1:50 - 1:100   |         |            |  |

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 284-563 of human MCCC2 (NP\_071415.1).

#### Synonyms

MCCB; MCCCbeta; MCCC2

WB

IHC-P

IF/ICC

## **Product Information**

€ www.abclonal.com

1:50 - 1:100

Isotype lgG

Purification Affinity purification


#### Storage

Source

Rabbit


Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

# **Validation Data**



Western blot analysis of extracts of various cell lines, using MCCC2 Rabbit pAb (A15181) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020).

Exposure time: 90s.



Immunofluorescence analysis of U2OS cells using MCCC2 antibody (A15181) at dilution of 1:100. Blue: DAPI for nuclear staining.