Leader in Biomolecular Solutions for Life Science

RXRα Rabbit pAb

Catalog No.: A15242



Basic Information

Observed MW

51kDa

Calculated MW

51kDa

Category

Mouse Monoclonal Antibody

Applications

WB, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

Retinoid X receptors (RXRs) and retinoic acid receptors (RARs) are nuclear receptors that mediate the biological effects of retinoids by their involvement in retinoic acid-mediated gene activation. These receptors function as transcription factors by binding as homodimers or heterodimers to specific sequences in the promoters of target genes. The protein encoded by this gene is a member of the steroid and thyroid hormone receptor superfamily of transcriptional regulators. Alternative splicing of this gene results in multiple transcript variants.

Recommended Dilutions

WB

1:500 - 1:2000

Immunogen Information

Gene ID 6256 **Swiss Prot**

P19793

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human RXR α (NP 002948.1).

Synonyms

NR2B1; RXRalpha; RXR-alpha; RXRα

Contact



www.abclonal.com

Product Information


Source Rabbit **Isotype** IgG **Purification**Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

Validation Data



Western blot analysis of extracts of various cell lines, using RXR α antibody (A15242) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 120s.