Leader in Biomolecular Solutions for Life Science

# MAPK6 Rabbit pAb

Catalog No.: A15304



### **Basic Information**

# **Observed MW**

120kDa

#### **Calculated MW**

83kDa

#### Category

Polyclonal Antibody

# **Applications**

WB,IF/ICC,ELISA

#### **Cross-Reactivity**

Human, Mouse, Rat

# **Background**

The protein encoded by this gene is a member of the Ser/Thr protein kinase family, and is most closely related to mitogen-activated protein kinases (MAP kinases). MAP kinases also known as extracellular signal-regulated kinases (ERKs), are activated through protein phosphorylation cascades and act as integration points for multiple biochemical signals. This kinase is localized in the nucleus, and has been reported to be activated in fibroblasts upon treatment with serum or phorbol esters.

# **Recommended Dilutions**

**WB** 1:200 - 1:2000

**IF/ICC** 1:50 - 1:200

# **Immunogen Information**

**Gene ID Swiss Prot** 5597 Q16659

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 372-721 of human MAPK6 (NP\_002739.1).

### **Synonyms**

ERK3; PRKM6; p97MAPK; HsT17250; MAPK6

### **Contact**

www.abclonal.com

# **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**




Western blot analysis of various lysates using MAPK6 Rabbit pAb (A15304) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 15s.



Immunofluorescence analysis of L929 cells using MAPK6 Rabbit pAb (A15304) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.