### A15372

Leader in Biomolecular Solutions for Life Science

# FIG4 Rabbit pAb

Catalog No.: A15372



#### **Basic Information**

Observed MW 104kDa

Calculated MW 104kDa

Category Polyclonal Antibody

Applications WB,ELISA

Cross-Reactivity Human,Mouse

#### Background

The protein encoded by this gene belongs to the SAC domain-containing protein gene family. The SAC domain, approximately 400 amino acids in length and consisting of seven conserved motifs, has been shown to possess phosphoinositide phosphatase activity. The yeast homolog, Sac1p, is involved in the regulation of various phosphoinositides, and affects diverse cellular functions such as actin cytoskeleton organization, Golgi function, and maintenance of vacuole morphology. Membrane-bound phosphoinositides function as signaling molecules and play a key role in vesicle trafficking in eukaryotic cells. Mutations in this gene have been associated with Charcot-Marie-Tooth disease, type 4J.

## **Recommended Dilutions**

#### **Immunogen Information**

WB

#### 1:200 - 1:2000

**Gene ID** 9896 Swiss Prot Q92562

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 608-907 of human FIG4 (NP\_055660.1).

#### Synonyms

BOP; YVS; BTOP; SAC3; ALS11; CMT4J; KIAA0274; dJ249I4.1; FIG4

Contact
---------

## **Product Information**


www.abclonal.com

**Source** Rabbit **Isotype** IgG Purification Affinity purification

#### Storage

Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

## Validation Data



Western blot analysis of various lysates using FIG4 Rabbit pAb (A15372) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution. Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit (RM00020). Exposure time: 30s.