# ABclonal www.abclonal.com

## **ZFX** Rabbit pAb

Catalog No.: A16444

## **Basic Information**

## **Observed MW**

110kDa

## **Calculated MW**

91kDa

## Category

Primary antibody

## **Applications**

ELISA,WB

#### **Cross-Reactivity**

Human, Mouse, Rat

## **Background**

This gene on the X chromosome is structurally similar to a related gene on the Y chromosome. It encodes a member of the krueppel C2H2-type zinc-finger protein family. The full-length protein contains an acidic transcriptional activation domain (AD), a nuclear localization sequence (NLS) and a DNA binding domain (DBD) consisting of 13 C2H2-type zinc fingers. Studies in mouse embryonic and adult hematopoietic stem cells showed that this gene was required as a transcriptional regulator for self-renewal of both stem cell types, but it was dispensable for growth and differentiation of their progeny. Multiple alternatively spliced transcript variants encoding different isoforms have been identified, but the full-length nature of some variants has not been determined.

## **Recommended Dilutions**

WB

1:500 - 1:2000

## Immunogen Information

Gene ID

**Swiss Prot** 

7543

P17010

## Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 180-380 of human ZFX (NP\_003401.2).

## **Synonyms**

ZNF926; ZFX

## **Contact**

•

www.abclonal.com

## **Product Information**

**Source** Rabbit Isotype


**Purification** 

IgG Affinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.



Western blot analysis of various lysates using ZFX Rabbit pAb (A16444) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021).

Exposure time: 90s.