

A1654

Leader in Biomolecular Solutions for Life Science



## PTHLH Rabbit pAb

Catalog No.: A1654

4 Publications

### Basic Information

#### Observed MW

20kDa

#### Calculated MW

20kDa

#### Category

Mouse Monoclonal Antibody

#### Applications

WB, ELISA

#### Cross-Reactivity

Human, Mouse, Rat

### Background

The protein encoded by this gene is a member of the parathyroid hormone family. This hormone, via its receptor, PTHR1, regulates endochondral bone development and epithelial-mesenchymal interactions during the formation of the mammary glands and teeth. It is responsible for most cases of humoral hypercalcemia of malignancy, and mutations in this gene are associated with brachydactyly type E2 (BDE2). Alternatively spliced transcript variants have been found for this gene. There is also evidence for alternative translation initiation from non-AUG (CUG and GUG) start sites, downstream of the initiator AUG codon, resulting in nuclear forms of this hormone.

### Recommended Dilutions

WB

1:100 - 1:500

### Immunogen Information

#### Gene ID

5744

#### Swiss Prot

P12272

#### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 25-175 of human PTHLH (NP\_945315.1).

#### Synonyms

HHM; PLP; BDE2; PTHR; PTHRP; PTHLH

### Contact



[www.abclonal.com](http://www.abclonal.com)

### Product Information

#### Source

Rabbit

#### Isotype

IgG

#### Purification

Affinity purification


#### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH 7.3.

## Validation Data

---



Western blot analysis of extracts of HepG2 cells, using PTHLH antibody (A1654) at 1:500 dilution.  
Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.  
Lysates/proteins: 25µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit (RM00020).  
Exposure time: 180s.