Leader in Biomolecular Solutions for Life Science

# **ZNF763 Rabbit pAb**

Catalog No.: A16615



# **Basic Information**

### **Observed MW**

46kDa

#### **Calculated MW**

46kDa

#### Category

Polyclonal Antibody

#### **Applications**

WB,ELISA

#### **Cross-Reactivity**

Human, Mouse

# **Background**

Predicted to enable DNA-binding transcription factor activity and RNA polymerase II cisregulatory region sequence-specific DNA binding activity. Predicted to be involved in regulation of transcription by RNA polymerase II. Predicted to be located in nucleus.

# **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

**Gene ID** 284390

Swiss Prot Q0D2J5

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 315-370 of human ZNF763 (NP\_001012771.1).

# **Synonyms**

ZNF; ZNF440L; ZNF763

# **Contact**

•

www.abclonal.com

# **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using ZNF763 Rabbit pAb (A16615) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (RM00021).

Exposure time: 10s.

Western blot analysis of various lysates using ZNF763 Rabbit pAb (A16615) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking-fuffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 30s.

