

A17114

Leader in Biomolecular Solutions for Life Science



TDP-43/TARDBP Rabbit pAb

Catalog No.: A17114

Basic Information

Observed MW

Refer to figures

Calculated MW

45kDa

Category

Mouse Monoclonal Antibody

Applications

WB, IHC-P, IF/ICC, ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

HIV-1, the causative agent of acquired immunodeficiency syndrome (AIDS), contains an RNA genome that produces a chromosomally integrated DNA during the replicative cycle. Activation of HIV-1 gene expression by the transactivator Tat is dependent on an RNA regulatory element (TAR) located downstream of the transcription initiation site. The protein encoded by this gene is a transcriptional repressor that binds to chromosomally integrated TAR DNA and represses HIV-1 transcription. In addition, this protein regulates alternate splicing of the CFTR gene. A similar pseudogene is present on chromosome 20.

Recommended Dilutions

WB	1:500 - 1:2000
IHC-P	1:50 - 1:200
IF/ICC	1:50 - 1:200

Immunogen Information

Gene ID

23435

Swiss Prot

Q13148

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-260 of human TDP-43/TARDBP (NP_031401.1).

Synonyms

ALS10; TDP-43; TDP-43/TARDBP

Contact



www.abclonal.com

Product Information

Source

Rabbit

Isotype

IgG

Purification


Affinity purification

Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.


Validation Data




Immunohistochemistry analysis of TDP-43/TARDB in paraffin-embedded rat spleen using TDP-43/TARDB Rabbit pAb (A17114) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of TDP-43/TARDB in paraffin-embedded human gastric cancer using TDP-43/TARDB Rabbit pAb (A17114) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of TDP-43/TARDB in paraffin-embedded mouse brain using TDP-43/TARDB Rabbit pAb (A17114) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of U-2 OS cells using TDP-43/TARDB Rabbit pAb (A17114) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.