Leader in Biomolecular Solutions for Life Science

# SYT12 Rabbit pAb

Catalog No.: A17241



# **Basic Information**

### **Observed MW**

46kDa

#### **Calculated MW**

47kDa

#### Category

Polyclonal Antibody

### **Applications**

WB,IF/ICC,ELISA

# **Cross-Reactivity**

Mouse

# **Background**

This gene is a member of the synaptotagmin gene family and encodes a protein similar to other family members that mediate calcium-dependent regulation of membrane trafficking in synaptic transmission. Studies of the orthologous gene in rat have shown that the encoded protein selectively modulates spontaneous synaptic-vesicle exocytosis and may also be involved in regulating calcium independent secretion in nonneuronal cells. Alternative splicing results in multiple transcript variants. The gene has previously been referred to as synaptotagmin XI but has been renamed synaptotagmin XII to be standard with mouse and rat official nomenclature.

# **Recommended Dilutions**

**WB** 1:500 - 1:2000

IF/ICC 1:50 - 1:200

# **Immunogen Information**

**Gene ID**91683

Swiss Prot
908|V01

#### **Immunogen**

Recombinant fusion protein containing a sequence corresponding to amino acids 95-200 of human SYT12 (NP\_001171351.1).

### **Synonyms**

SYT11; sytXII; SYT12

### **Contact**

www.abclonal.com

# **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**




Western blot analysis of lysates from Mouse brain, using SYT12 Rabbit pAb (A17241) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 3min.



Immunofluorescence analysis of L929 cells using SYT12 Rabbit pAb (A17241) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.