Leader in Biomolecular Solutions for Life Science

# **CCR4 Rabbit pAb**

Catalog No.: A17324



### **Basic Information**

### **Observed MW**

70kDa

#### **Calculated MW**

41kDa

#### Category

Polyclonal Antibody

# **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human

# **Background**

The protein encoded by this gene belongs to the G-protein-coupled receptor family . It is a receptor for the CC chemokine - MIP-1, RANTES, TARC and MCP-1. Chemokines are a group of small polypeptide, structurally related molecules that regulate cell trafficking of various types of leukocytes. The chemokines also play fundamental roles in the development, homeostasis, and function of the immune system, and they have effects on cells of the central nervous system as well as on endothelial cells involved in angiogenesis or angiostasis.

# **Recommended Dilutions**

WB

1:500 - 1:2000

# **Immunogen Information**

Gene ID 1233 Swiss Prot P51679

#### **Immunogen**

A synthetic peptide corresponding to a sequence within amino acids 1-100 of human CCR4 (NP 005499.1).

### **Synonyms**

CKR4; K5-5; CD194; CMKBR4; ChemR13; CC-CKR-4; HGCN:14099; CCR4

### **Contact**

•

www.abclonal.com

# **Product Information**


**Source** Rabbit **Isotype** IgG **Purification**Affinity purification

# Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.01% thimerosal,50% glycerol,pH7.3.

# **Validation Data**



Western blot analysis of various lysates using CCR4 Rabbit pAb (A17324) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane. Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 120s.